

Calculation of concentration fields of high-inertia aerosol particles in the flow past a cylindrical fibre

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© Published under licence by IOP Publishing Ltd. The behaviour of high-inertia aerosol particles' concentration fields in stationary gas suspension flows around a cylinder is investigated using a numerical solution to the Navier-Stokes equations and the fully Lagrangian approach for four Stokes numbers ($Stk = 0.1, 1, 4, 10$) and three Reynolds numbers ($Re = 1, 10, 100$). It has been shown that the points of maximum particle concentration along each trajectory shift downstream both when Stk and/or Re increase.

<http://dx.doi.org/10.1088/1755-1315/107/1/012109>

References

- [1] Crowe C T 1982 Trans. ASME J. Fluids Eng. 104 303
- [2] Ferry J and Balachandar S 2001 International Journal of Multiphase Flow 27 226
- [3] Desjardins O, Fox R O and Villedieu P 2008 Journal of Computational Physics 227 39
- [4] Healy D P and Young J B 2005 Proc. R. Soc. A. 461 225
- [5] Fernandez de la Mora J and Rosner D E 1981 Physicochem. Hydrodyn. 2 21
- [6] Osipov A N 2000 Astrophysics and Space Science 274 86
- [7] Zaripov T S, Gilfanov A K, Begg S M, Rybdylova O D, Sazhin S S and Heikal M 2017 Atomization and Sprays 27 510